

Combining Trig Functions and Inverse Trig Functions - Part I



Preliminaries:

- Trig functions
- Inverse Trig Functions

Objectives:


- Find values when trig functions and inverse trig functions are combined.

Example 1

Find

$$\sin\left(\cos^{-1}\left(\frac{1}{2}\right)\right)$$

↑
number




Example 1

Find

$$\sin\left(\cos^{-1}\left(\frac{1}{2}\right)\right)$$

↑
angle




$$\theta = \cos^{-1}\left(\frac{1}{2}\right)$$

Example 1

Find

$$\sin \left(\cos^{-1} \frac{1}{2} \right)$$


number



Example 1

Find

$$\sin \left(\cos^{-1} \frac{1}{2} \right) = \sin \left(\frac{\pi}{3} \right) = \frac{\sqrt{3}}{2}$$




Example 2

Find

$$\sin \left(\cos^{-1} \frac{2}{3} \right)$$

number




Example 2

Find

$$\sin \left(\cos^{-1} \frac{2}{3} \right)$$

angle




$$\theta = \cos^{-1} \left(\frac{2}{3} \right)$$

Example 2

Find


$$\sin\left(\cos^{-1}\frac{2}{3}\right)$$

number



Example 2

If $\cos \theta = \frac{2}{3}$ and θ is in quadrant I, find $\sin \theta$.



Example 2


$$\left(\frac{2}{3}\right)^2 + (\sin \theta)^2 = 1$$

$$(\sin \theta)^2 = 1 - \frac{4}{9}$$

$$(\sin \theta)^2 = \frac{5}{9}$$

$$\sin \theta = \frac{\sqrt{5}}{3}$$

$$\sin\left(\cos^{-1}\frac{2}{3}\right) = \frac{\sqrt{5}}{3}$$



Example 2

$$\sin\left(\cos^{-1}\frac{2}{3}\right)$$


$$b^2 + 2^2 = 3^2$$

$$b^2 = 9 - 4 = 5$$

$$b = \pm\sqrt{5}$$

$$\sin \theta = \frac{b}{3} = \frac{\sqrt{5}}{3}$$


$$\sin\left(\cos^{-1}\frac{2}{3}\right) = \frac{\sqrt{5}}{3}$$



Example 3 - Negative Values

Find


$$\sin \left(\cos^{-1} -\frac{2}{3} \right)$$
$$= \sin \theta = \frac{\sqrt{5}}{3}$$



Example 4

Find


$$\sin \left(\sin^{-1} .276 \right)$$
$$\sin \theta = .276$$



Example 5

Find


$$\cos^{-1} \left(\sin \frac{\pi}{3} \right)$$
$$= \cos^{-1} \left(\frac{\sqrt{3}}{2} \right)$$
$$= \frac{\pi}{6}$$



Example 5

Find $\cos \left(\tan^{-1} 2 \right)$

$$\tan \theta = \frac{2}{1}$$
$$\cos \theta = \frac{1}{\sqrt{5}} = \frac{\sqrt{5}}{5}$$
$$\cos \left(\tan^{-1} 2 \right) = \frac{\sqrt{5}}{5}$$




Example 6

$$\text{Find } \tan\left(\cos^{-1}\frac{3}{5}\right)$$

$$\theta = \cos^{-1}\left(\frac{3}{5}\right)$$

$$\tan\theta = \frac{4}{3}$$

$$\tan\left(\cos^{-1}\frac{3}{5}\right) = \frac{4}{3}$$



Example 7


$$\text{Find } \sec\left(\sin^{-1}\frac{1}{4}\right)$$

$$\theta = \sin^{-1}\frac{1}{4}$$

$$\cos\theta = \frac{\sqrt{15}}{4}$$

$$\sec\theta = \frac{4}{\sqrt{15}} = \frac{4\sqrt{15}}{15}$$

$$\sec\left(\sin^{-1}\frac{1}{4}\right) = \frac{4\sqrt{15}}{15}$$



Example 8


$$\tan\left(\sec^{-1}\frac{5}{2}\right)$$

$$= \tan\left(\cos^{-1}\frac{2}{5}\right)$$

$$\theta = \cos^{-1}\frac{2}{5}$$

$$\tan\theta = \frac{\sqrt{21}}{2}$$

$$\tan\left(\sec^{-1}\frac{5}{2}\right) = \frac{\sqrt{21}}{2}$$



Recap

- Inputs to inverse trig functions give information about the lengths of sides of a triangle.
- Label a triangle using this information.
- Find the missing side by the Pythagorean Theorem
- Read the values of the other trig functions from the triangle