# **Graphing Rational Functions**



University of Minnesota Graphing Rational Functions

## **Preliminaries and Objectives**

#### Preliminaries

- Intercepts
- Factoring Polynomials
- Graphing Polynomials
- Long Division of Polynomials

Objectives

Graph Rational Functions

$$f(x) = \frac{x^5 + 5x^4 - 5x^3 - 45x^2 + 108}{x^3 - x^2 - x + 1} = \frac{(x+3)^3(x-2)^2}{(x+1)(x-1)^2}$$

- y-intercept
- x-intercepts
- vertical asymptotes
- end behavior

Step 1: Find the *y*-intercept

$$f(x) = \frac{(x+3)^3(x-2)^2}{(x+1)(x-1)^2}$$

$$f(0) = \frac{(0+3)^3(0-2)^2}{(0+1)(0-1)^2} = 108$$

Step 2: Find the *x*-intercepts

$$f(x) = \frac{(x+3)^3(x-2)^2}{(x+1)(x-1)^2}$$

$$(x+3) = 0$$
 or  $(x-2) = 0$   
 $x = -3$  or  $x = 2$ 

#### x-intercepts at -3 and 2


Step 3: Find asymptotes

$$f(x) = \frac{(x+3)^3(x-2)^2}{(x+1)(x-1)^2}$$


f(x) is undefined when x = -1 and when x = 1

Step 4: Analyze intervals

$$f(x) = \frac{(x+3)^3(x-2)^2}{(x+1)(x-1)^2}$$




**Example 1** 




Example 2

$$f(x) = \frac{x^3 - x^2 - 8x + 12}{x^4 + 2x^3 - 2x - 1}$$



**Example 1** 



$$f(x) = \frac{x^2}{x^4 - 2x^2 + 1}$$

• If the degree of the numerator is less than the degree of the denominator, then

$$\lim_{x\to\infty} f(x) = 0 \text{ and } \lim_{x\to-\infty} f(x) = 0$$

$$f(x) = \frac{2x^2 - 18}{x^2 + 1}$$

 If the degree of the numerator is the same as the degree of the denominator, then long division will produce a constant and a remainder. In this case, the constant is a horizontal asymptote.


$$f(x) = \frac{3x^3 + x^2 + 3x + 6}{x^2 + 1}$$


 If the degree of the numerator is larger than the degree of the denominator, then long division will produce two terms and a remainder. In this case the graph approaches an oblique asymptote.


$$f(x) = \frac{3x^4 + x^3 - 3x^2 - x + 5}{x^2 - 1}$$


 If the degree of the numerator is at least two greater than the degree of the denominator, then the graph resembles x<sup>2</sup>, x<sup>3</sup>, x<sup>4</sup>,...


$$f(x) = \frac{x^2}{x^4 - 2x^2 + 1} = \frac{x^2}{(x - 1)^2(x + 1)^2}$$

















- Find *y*-intercept
- Find *x*-intercepts
- Find vertical asymptotoes
- Determine end behavior
- Sketch the graph